Stomach

11 January 2022 17:44

STOMACH


Fundus
F
C
Cardia
B
Body
PA
Pyloric Antrum

Cell type	Location	Function
1. Parietal cells	Body	HCl + Castel Intrinsic factor — B12 absorption
2. Mucus	Body & Antrum	mucus + HCO3⁻
3. Chief cells	Body	Pepsin
4. E.C.L cells	Body	Histamine
5. G-cells	Antrum	Gastrin
6. D-cells	Body & Antrum	Somatostatin

Acid physiology


Empty Stomach
⊕
D-cells
Somatostatin
⊘
G-cells
Gastrin
⊕
Parietal cells
↑HCl
CIF
Histamine
ECL cells
Acetylcholine
Ach
⊕
Vagus
MEAL
⊕ VAGUS

VAGUS


R
L
Hepatic branch
Sero muscular br.
Criminal Nerve of Grassi
Terminal Nerve of LATARJET
Celiac br.
Sero muscular br.

Blood Supply


Short Gastric Vessels
Splenic a
HA proper
LGA
CHA
Left gastric Epiploic a
Gastro duodenal Artery
m/c source for Bleeding in P.U.D
Right gastric Epiploic artery
Posterior superior pancreatico duodenal artery
Anterior superior pancreatico duodenal artery

CHPS : Congenital Hypertrophic Pyloric Stenosis

Since Birth → ↑ size of Circular muscles → Pyloric region → Narrowing → Gastric outlet obstruction.

C/F : Age : 3rd – 6th wk (4th wk)
Sex : M > F (4 : 1)
1st BORN MALE CHILD : ↑R/F

Pathogenesis

Ⓡ Episodes of vomiting → Non Bilious vomiting / Projectile vomiting / Extensive cry just before vomiting



O/E
- olive lump : Best noted after feeding
- visible peristalysis
- Starvation Stools : only mucus
- Dehydrated

S. Electrolyte : H2O + HCl + Na⁺
- H⁺ → m. Alkalosis
- Cl⁻ → Hypochloremia
- Na⁺ → ⊕ Aldosterone
- Hypokalemia
- Paradoxical Aciduria

A : Absorption of Na⁺ & H2O
B : BP maintains
C, D : CD of nephron → Na⁺ K⁺ / Na⁺ H⁺
E : Excretion of K⁺

Na⁺ K⁺ ✓
Na⁺ H⁺ ✓

1st INV : X-Ray : massive fundus gas.
JOC : USG : Double Track app



→ CHPS.

Q. **Mx : Sx : Lap. Ramsteds pyloromyotomy**

DUODENAL ATRESIA

Incomplete or Improper chanalization b/w Fore gut & mid gut

FORE GUT — D2
MID GUT — D3 to T2/3 Colon.
HIND GUT — Dentate line.

C/F : Age : Day 1 / Day 2
Sex : M = F



- Ⓡ Episodes of vomiting ≥ 30min of feeding
- Bilious vomiting
- Non projectile

INV : X-Ray → Double Bubble Sign : Duodenal Atresia
→ Tripple Bubble Sign : Jejunal Atresia

A/w other Congenital Anomalies
- Annular pancreas
- Down Syndrome
- Extra hepatic Biliary Atresia

Mx : Exploratory laprotomy → Duodenoduodenostomy



Peptic Ulcer Disease : P.U.D

A/w Gastric type of mucosa — Breach in Continuity of mucus membrane.

Duodenum (D1) (m/c) | Stomach | Lower 1/3 Esophagus | Meckel's Diverticulum.

modified Johnson's Classn



Type I : Ulcer @ lesser curvature — Angularis Incisura.
Type II : 2 ulcers — Body, D1
Type III : Pre pyloric gastric ulcer
Type IV : High on lesser curvature.
Type V : N.S.A.I.D.S induced Gastric Ulcer.

Ⓒ Peptic Ulcer Disease

- H. Pylori → 85-90% Duodenal Ulcer; 70% Gastric ulcer
- Smoking
- NSAIDS
- Steroids
- Stress ulcers → Burns : Curling ulcer; Head injury : Cushing ulcer
- Endocrine Ⓒ : - Gastrinoma (Zollinger Ellison Synd) - Cushing Syndrome - Hyper para thyroidism

Pathophysiology

	Duodenal ulcer	Gastric ulcer : PRE-malignant
m/c/s :	D1	Angularis Incisura
Status :	Hyper acidic	Normo acidic / Hypoacidic
Pain :	≥ 2hrs of Eating	Immediately after eating — Food pain
Intensity :	mild to mod.	Severe
Appetite :	↑↑	↓↓
Weight :	Gain	loss
Bleeding :	-	m/c → posterior Gastric ulcer — m/c source : Gastroduodenal artery
Perforation :	m/c → Anterior ulcers perforate.	-

JOC : U.G.I.E

Gastric ulcer → Biopsy from all quadrants → No Cancer / Adeno Ca.
No Cancer → H. Pylori Eradication
Tripple therapy : PPI + Broad spectrum AB + Tinidazole
Quadraple therapy : Tripple + Bismuth / Secnidazole] 10-14 days
→ Repeat Endoscopy → Ulcer → Healed : observation & follow up c̄ annual UGIE / persistent : Ⓡ Biopsy → No Cancer / Adeno Ca → Surgical Excision → Billroth-I / Billroth-II / Antrectomy


Gastroduodenostomy
Gastrojejunostomy

Duodenal ulcer → H. Pylori Eradication → fails → Surgery → Truncal vagotomy / Selective vagotomy / Highly selective vagotomy


TRUNCAL
SELECTIVE
Highly selective vagotomy

SURGERY : ↓ maximum acid out put — TRUNCAL VAGOTOMY + Antrectomy.

Complications of P.U.D

- Bleeding (Hemorrhage) → Gastric ulcers — posteriorly situated ulcer — m/c source : Gastro duodenal a
- Upper G.I Bleeding

JOC : UGIE : FORREST CLASSIFICATION



Ia : Active pulsatile Bleeding Visible Blood vessel
Ib : Oozing of Blood Visible Blood vessel
IIa : Non Bleeding Visible Blood vessel
] High Risk for Recurrance.
IIb : Adherent clot] Intermediate R/F Recurrance.
IIc : Black Dot
III : Clean Base ulcer
] low risk for Recurrance.

Mx : Active Bleeding : Hemo clips



THREE POINT LIGATION of Gastro duodenal artery

- Perforation (Hollow viscus Perforation)
  - m/c : Duodenal ulcer
  - Anteriorly situated ulcer

C/F : Perforation Peritonitis



Contamination
1. Bile : most Irritant for Peritoneum.
2. Food substance
3. Blood
4. G.I secretion

- Severe abdominal pain
- High grade fever
- Nausea & vomiting — paralytic ileus.
- Non passage of Stools & flatus.
- Abdominal Distention.

O/E :
- Guarding ⊕
- Cardboard Rigidity ⊕
- Bowel Sounds ⊖
- ↑HR, ↑RR, Hypotension

JOC : Erect X-Ray Abdomen → ⊕ free gas under Right Hemi diaphragm

BEST : CECT abdomen

Mx : Exploratory laprotomy
Gastric ulcer → Biopsy from the Edge → ⊕ Adeno carcinoma → Mx as Gastric Carcinoma / Benign ulcer
Duodenal ulcer → Graham's omental patching

- Gastric outlet obstruction : m/c D1



C/F
- Ⓡ Episodes of Non Bilious vomiting
- Colicky Abdominal pain

1st INV : X-Ray
JOC : UGIE
Mx : ✓ Heineke-mikulicz Pyloroplasty
- Finney's pyloroplasty




TYPE-II
TEA POT STOMACH
HOUR GLASS STOMACH : Type-I

Complications a/w P.U.D SURGERY

BILROTH Sx : II → Duodenal Stump Blow out : 4th P.O.D / Pernicious anemia (B12 L 2° to ↓CIF) / Dumping Syndrome
VAGOTOMY → Truncal vagotomy → Hyposecretion / Cholelithiasis / Gastric paresis → Drainage procedure





Dumping Syndrome → Early / Late

Early : Ⓒ Rapid transit of food from stomach into jejunum
Late : Ⓒ Rapid transit of food from stomach into jejunum


sugar Rich
Draw water into jejunum
Osmotic Diarrhoea

DOC : Octreotide

Sugar Rich Bolus in jejunum → ↑↑↑ Insulin → Severe Hypoglycemia

	Early	Late
Time of Presentation :	< 20-30 mins	> 2hr
C/F :	Diarrhoea	Hypoglycemia
Precipitating factor :	Food	Exercise
Relieving factor :	lie down supine	Food

Upper G.I Bleed

DIELAFOY'S GASTRIC LESION

- Abnormal Arteriole of ~3mm Ø located 6-10cm from G.E.J on the lesser Curvature of Stomach.



C/F
- Hematemesis Ⓡ
- malena ⊕
- Hypotension

JOC : UGIE
Mx : Electro coagulation of the a/w arteriole.

G.A.V.E : Gastric Antral vascular Ectasia — WATER MELON STOMACH

- Dilated vessels in the antrum of the Stomach.


Water melon Stomach
Dilated Vessels.

C/F
- AGE > 60yrs
- SEX : F > M
- Anemia : Chronic Blood loss.

JOC : UGIE
Mx : UGIE → Electrocoagulation.

MENETRIERS DISEASE

- Hypertrophied mucosal folds in the Body and fundus of Stomach resulting in Hypoalbuminemia (Protein losing) Hypo-


mucus producing cells
Surface
Isthmus
Neck
Parietal cells : HCl CIF
ECL cells
Base
Chief cells.
GASTRIC PIT

Ⓒ → Idiopathic > Strong association → Adult : H. Pylori / Children : CMV

C/F : AGE : 40 - 50yrs
SEX : F > M

- Diarrhoea
- Hypochlorhydria / Achlorhydria
- Pernicious Anemia : PRE-MALIGNANT
- Hypergastrinemia
- Hypoproteinemia : most charch. feature.
- Steatorrhea

JOC : UGIE
Mx : Prophylactic Total Gastrectomy.

Carcinoma Stomach

m/c/s : Around the Gastro Esophageal junction


19% Ca
15% Ca
C
Body
Antrum
Pylorus
7%
18%
60% of Total malignancy

m/c/s c̄ in the : Cardia Stomach

R/F
- NUTRITIONAL R/F : - ↓ Fat diet - ↓ protein diet - Salted meat - High Nitrate Consumption
- Environmental R/F : - Poor preparation of food : Smoked - Smoking - Poor drinking water - lack of refrigeration.
- medical R/F : - Prior gastric Sx - H. Pylori : (Cag-A +ve) Strongest R/F Ca - Gastric atrophy - Type A / B gastritis (Autoimmune / H. Pylori) - Adenomatous polyps. - Males > Females

Gastric Ca Genetics

1. E-CADHERIN : ↑ 80%
2. F.A.P
3. Duodenal polyps : Very High malignant potential
4. p53 mutation
5. HNPCC
6. p16

WHO classn : Carcinoma Stomach

Adeno Ca (m/c) → Papillary : Good / Tubular / Mucinous / Signet ring : WORST prognosis / mixed
Adeno Squamous | Squamous Ca | Undiff | Unclassified

LAUREN Classn : Gastric Ca

	Intestinal Type	Diffuse Type
SEX	M	F
AGE	> 50yrs / Ml+2+Age	Young (30-40yrs)
Genetic	Sporadic	Familial
Genes	APC, P53, P16	E-cadherin, P53, P16
R/F	Gastric Atrophy, Intestinal metaplasia	Blood group 'A'
mode of spread	Blood	Direct & lymph
H.P.E	- Gland forming - MSI	Signet ring
Prognosis	Good	Poor

C/F : Silent Ca (slow growth)
- Dysphagia : Solids > Liq, progressive
- Dyspepsia ⊕
- Early satiety
- Gastric outlet obstruction.
- Hematemesis. - malena. - Anemia.
- Vague Epigastric discomfort



Q. m/c clinical presentation : Abdominal Pain > Weight loss.

ATYPICAL PRESENTATION

1. Enlarged lymph nodes → Ⓛ Supra clavicular : Virchow LN (Troisier Sign); Ⓛ Axilla : Irish LN; Periumbilical : Sister mary Joseph LN
2. Migratory Superficial Thrombophlebitis : (Trousseau Syndrome)
3. Pelvic metastasis : Per rectal → ⊕ Blummer Self.
4. Ovary tumour : 2° to Drop metastasis — KRUKENBERG TUMOUR / Trans peritoneal Spread

JOC : UGIE + Bx

Morphology

Q. Early Ca. — Japanese classn | Advanced Ca — Borrman's classn



Type I : PROTRUDED
Type IIa : Flat & Sup. Elevated
Type IIb : Flat
Type IIc : Depressed
Type III : Excavated

Type I : Polypoidal
Type II : Ulcer c̄ Elevated margin.
Type III : Ulcer c̄ infiltration into gastric wall.
Type IV : Linitis plastica : a/w worst prognosis
Type V : Non classifiable.

TNM - Classification

T1a : ⊕ lamina Propria
T1b : ⊕ Sub mucosa
T2 : ⊕ muscularis propria
T3 : ⊕ Sub serosa
T4a : Tumour perforates Serosa
T4b : Tumour invading adjacent organs.

N0
N1 : 1 - 2 regional L.N ⊕
N2 : 3 - 6 " "
N3a : 7 - 15 " "
N3b : > 15 " "

M0 : ✗
M1 : ✓

Mx

Operable Tumour → ≤ T3, N1, N2
Inoperable Tumour → ≥ T4 or N3 or M1 → palliative mx → - Feeding jejunostomy - Endo G.I Stenting


proximal 1/3 : Extended gastrectomy + D2
middle 1/3 : Total gastrectomy + D2
Distal 1/3 : Sub-Total gastrectomy + D2
Sx

+ Chemo : E : Epirubicin, C : Cisplatin, F : 5FU

Gastro Intestinal Stromal Tumour

G.I.S.T

- m/c mesenchymal tumour of G.I.T
- m/c/s for G.I.S.T : Stomach

Genetics of G.I.S.T

Tumour markers:
- over Expression of Tyrosine kinase Receptor (C-kit)
- CD117
- CD34
- BCL-2

New Tumour marker
- DOG-1

Origin : Interstitial cells of CAJAL

C/F
- Upper G.I Bleeding — m/c clinical presentation of G.I.S.T
- m/c/s for 2° : Liver
- Lymph nodes are usually spared in G.I.S.T

CARNEY'S TRIAD → Pulmonary Chondroma / Extra Adrenal paraganglioma / multi focal G.I.S.T

1st INV : UGIE
JOC : CECT

FLETCHER'S Classn

Benign:

	Size	Mitosis / 50HPF
1. Benign	< 2cm	< 5
2. Probably Benign	2-5cm	< 5

malignant:

	Size	Mitosis / 50HPF
low	< 2cm	> 5
mod.	2-5cm	> 5
	< 10cm	< 5
High	> 5cm	> 5

Mx → Benign → Wedge shaped Excision.
malignant → Resectable Tumour → Resection & anastamosis (margin : 5cm)
→ Unresectable Tumour → Imatinib → RESISTANCE → SUNITINIB

✓ Joc for Assessment of Recurrance : PET- Scan.

Polyps

- Inflammatory — (pseudo polyps) — (2° to U.C) — Low R/F malig. Transformation
- Hyperplastic polyp
- Adenomatous polyp — max. malig. Potential
- Hamartomatous polyp — Juvenile Polyps : Young Pt

PEUTZ-JEGHER Syndrome : GENE : STK-11 / LKB-1
- Polyposis (multiple polyps) + Pigmentation (muco cutaneous)
- m/c/s for Polyps : Jejunum
- m/c type of polyp : Hamartomatous Polyp.
- ↑ R/F Ca : PANCREATIC Ca, PERI AMPULLARY Ca, COLORECTAL Ca, THYROID Ca, DUODENAL Adeno Ca

Adenomatous Polyps → Tubular polyp / Tubulovillous polyp / Villous polyp — max R/F malignancy.
→ Pedunculated polyp / Sessile polyp


Apex
Head
Neck
Stalk
Base

HAGGIT CLASSIFICATION


LEVEL : 0 APEX
LEVEL : I HEAD
LEVEL : II NECK
LEVEL : III STALK
LEVEL : IV BASE

Endoscopic polypectomy — Level 0–III
Adeno ca of Colon. — Level IV

Polyposis

multiple polyps.

Familial Adenomatous Polyposis : FAP

- Chr 5
- Gene : APC
- Polyposis : min. no. of polyps req > 100
  Attenuated FAP : < 100 polyps
- A.D
- 100% R/F malignant transformation.

Pt. c̄ FAP

- Total proctocolectomy c̄ ileo-Anal Anastamosis (J-pouch)

Screen all family members
R/F malignancy ↑↑ > 40yrs
→ - Colonoscopy - mutation → ⊕ Prophylactic Total proctocolectomy / ⊖

Hereditary Non polyposis Colorectal Ca

H.N.P.C.C

- Genetic mutation : DNA mismatch Repair Gene
  Gene : MSH2 / MLH-1 : m/c

LYNCH → LYNCH-1 : ↑R/F : Colorectal Ca; LYNCH-2 : ↑R/F : - Uterine Ca - Ovarian Ca - Gastric Ca

HNPCC : Amsterdam Criteria (modified)

1. Rule out F.A.P
2. Atleast 3 family members ⊕
3. Atleast 2 Successive generation ⊕
4. Atleast 1 1st Degree relative ⊕
5. Atleast 1 Should be diagnosed < 50yrs.

GARDNER SYNDROME

1. FAP
2. Osteoma : maxilla / mandible
3. Fibroma
4. Epidermal Cyst
5. Desmoid tumour
6. Congenital Hypertrophy of Retinal pigment Epithelium

TURCOT Syndrome

- FAP + Brain tumours

COLORECTAL - Cancer

m/c type : Adeno Ca
m/c/s : Recto sigmoid.

R/F
- High fat intake c̄ low fiber diet
- Red meat
- I.B.D (UC > CD)
- Diverticular disease
- Adenomatous polyp
- FAP, HNPCC, Turcot Synd, Gardner Synd
- Ureterosigmoid anastamosis
- H/o pelvic radiation

Colorectal Ca

Ulcerative Type | Cauliflower Type | Stenosing Type | Tubular Type


m/c Right
m/c : Left
R
L
T2/3
T1/3

C/F

Right sided Ca	Left sided Ca
- malena] m/c - Anemia - Hematochezia : Rare - R.I.F mass : palpable - S/s of Small Bowel #	- large Bowel dynamic obstruction - Alteration in Bowel habits - Early morning spurious diarrhoea - Tenesmus - ⊕ Blood & mucus in stools - Hematochezia

JOC : Colonoscopic Biopsy
JOC for Staging : PET-CT
JOC for 'T' & 'N' Determination : MRI c̄ Endorectal coil

TNM - classn

T1 : ⊕ Submucosa
T2 : ⊕ muscular layer
T3 : ⊕ perimuscular connective tissue
T4a : > serosa (perforated)
T4b : ⊕ adjacent organ

N1 : 1-3 Regional L.N ⊕
N2 : ≥ 4 " "

M0 : ✗
M1 : ✓

DUKES STAGING

A : Limited to mucosa & submucosa
B1 : B/w muscular layer & serosa
B2 : Serosal perforation
C1 : B1 + L.N
C2 : B2 + L.N
D : Distant metastasis : m/c route of spread : Blood; m/c site for 2° : liver


- Small
- peripheral
- multiple
- hypovascular.

Mx : Margin : 5cm.


Extended Ⓡ Hemi Colectomy
Extended Ⓛ Hemi-Colectomy
Ⓛ Hemi Colectomy
Ⓡ Hemi Colectomy
R
L
MCA
RCA
LCA
JCA
SA

Right Hemi Colectomy "Radical"


R
L
MCA
RCA
JCA
10-12cm of T. ileum

Ileo transverse anastamosis

Extended Ⓡ Hemi colectomy "Radical"


Water shead area : Griffith point
R
L
MCA
RCA
JCA
12cm
Sudeck point

Left Hemi Colectomy Radical


L
MCA
LCA
SA
Beginning of rectum.

Extended Ⓛ Hemi Colectomy "Radical"


MCA
LCA
IMA
SA

Mx : Rectal Ca



Proximal margin : 5cm
Distal margin : 2cm.

Criteria for local Resection in Rectal ca

- T1N0 or T2N0
- < 4cm in diameter
- < 40% of Circumference of rectum
- < 10cm from dentate line
- well to mod. diff tumour.
- lympho vascular spread ⊖

Distal margin → Adequate / Inadequate

Adequate:



→ Primary Resection and anastamosis → Low anterior Resection / T.A.H.S (Trans anal mesorectal Excision Sx) / T.M.E Total meso rectal Excision.

Inadequate → A.P.R (Abdomino perineal) Resection

CHEMOTHERAPY

- ⊕ Lymph Node metastasis
- T3 / T4 ↓

FOLFOX : 5 FU + OXALIPLATIN + Folinic acid
FOLFIRI : 5 FU + Folinic acid + IRINOTECAN
CAPECITABINE + OXALIPLATIN.

Radio Rx : only for Rectal ca.