




FMGE June '26

RECALL SESSION

**A Mega Recall Session By
India's Top Educators**

Q. Identify the marked structured in the image given below.

- a. Maxillary sinus
- b. Posterior ethmoidal sinus
- c. Sphenoidal sinus
- d. Anterior ethmoidal sinus





Q. A 48-year-old diabetic man presents with unilateral nasal obstruction and nasal polyps. CT of the paranasal sinuses shows the double-density sign. What is the most likely diagnosis?

- a. Invasive fungal sinusitis
- b. Allergic fungal sinusitis
- c. Chronic rhinosinusitis with polyps
- d. Maxillary sinus carcinoma

Q. A patient presents with itching of the ear and ear discharge. Image of the otoscopic examination of the external auditory canal is given. What is the most likely diagnosis?

- a. Acute otitis externa
- b. Otomycosis
- c. Chronic suppurative otitis media
- d. Keratosis obturans





Q. A patient develops difficulty singing high-pitched notes after undergoing thyroid surgery. Injury to which nerve is the most likely cause?

- a. Recurrent laryngeal nerve
- b. External branch of the superior laryngeal nerve
- c. Internal branch of the superior laryngeal nerve
- d. Hypoglossal nerve

Q. A 7-year-old child presents with fever, ear pain, and swelling behind the ear. The clinical image shows postauricular swelling. What is the most likely diagnosis?

- a. Perichondritis of the pinna
- b. Acute mastoiditis
- c. Malignant otitis externa
- d. Preauricular abscess






Q. A 14-year-old boy presents with intermittent mucoid discharge from a small opening along the anterior border of the lower third of the sternocleidomastoid muscle since childhood. There is no history of trauma or surgery. What is the most likely diagnosis?

- a. Thyroglossal fistula
- b. Bezold abscess
- c. Branchial fistula
- d. Dermoid sinus




Q. A 56-year-old man with poorly controlled diabetes presents with facial pain, nasal obstruction, periorbital swelling, and black nasal discharge for 5 days. MRI of the paranasal sinuses is given below. What is the most appropriate initial treatment?

- a. Vancomycin
- b. Oral Voriconazole
- c. IV Amphotericin B with debridement
- d. IV Fluconazole



Q. A video laryngoscopy of a patient with hoarseness reveals unilateral abductor vocal cord palsy with the affected vocal cord fixed in the paramedian position. Which type of thyroplasty is the most appropriate surgical treatment?

- a. Type I thyroplasty
- b. Type II thyroplasty
- c. Type III thyroplasty
- d. Type IV thyroplasty



ENT



THANK YOU